

Can radiative correction cause large neutrino mixing?

Gautam Dutta*

Harish Chandra Research Institute, Jhusi, Allahabad 211019, India

Abstract

We investigate whether radiative corrections can be responsible for the generation of large mixing in a pair of degenerate neutrino with same CP parity. We find that this mechanism is fine tuned and doesn't work for arbitrary mixing at the high scale.

The data from the solar neutrino and the atmospheric neutrino experiments can be explained through flavor oscillations of massive neutrinos. The favoured solution of both these problems are a pair of neutrinos with very small mass squared difference and large mixing [1,2]. The mass square difference required for the solar neutrino oscillation is less than $10^{-5}eV^2$ while for the atmospheric neutrinos it is $10^{-3}eV^2$. If neutrinos form the hot Dark Matter then the masses of all the neutrinos have to be of the order of 1eV. This, along with the very small mass square differences suggests that neutrino mass spectrum is almost degenerate. If neutrinos have Majorana mass, then one can have a pair of degenerate neutrinos with opposite CP parities. These are called Pseudo Dirac neutrinos. The mixing between such pairs can be shown to be maximal [3]. The smallness of neutrino mass can be obtained through the seesaw mechanism [4] in which neutrinos can have both Dirac and Majorana masses. Even in this mechanism Pseudo Dirac neutrinos with maximal mixing are possible for a suitable choice of the structure of right handed Majorana mass matrix [5]. While these models can generate only one maximal mixing in 3 generation scenario, the solar neutrino problem and the atmospheric neutrino problem requires two pairs with maximal mixing. Another problem is that for degenerate Majorana neutrinos, large mixing may not be stable under small radiative corrections. A pair of Majorana neutrinos with opposite CP parities is stable but for a pair with same CP parity large mixing collapse to 0 due to radiative correction [6]. A detailed analysis on the stability of the neutrino mass matrix and mixing for various type of mass hierarchy is done in [7–9]. Some authors [10] suggested that the same instability can cause large mixing at the low scale from an arbitrary mixing at the high scale. This mechanism is similar to the MSW mechanism where

*email: gautam@mri.ernet.in

the mixing angle gets enhanced at the resonance. As this requires the present scale of the universe to be at the ‘resonance’ as far as neutrino mixing is concerned, this is a fine tuned solution. This is because 0 mixing is a fixed point of the renormalisation group evolution of neutrino masses and mixing whereas maximal mixing is not [8]. The fact that for degenerate mass structure, large mixing is unstable have been discussed by other authors [7–9,11] We investigate whether such a mechanism can be at work to generate the required large mixing considering the range of mass square differences from the solar and atmospheric neutrino experiments.

Let m_1 and m_2 be the mass eigenvalues of two neutrinos at a high scale Λ . Let θ be the mixing angle at this scale. Then in the flavour basis the mass matrix will be given as

$$M_F(\Lambda) = \begin{pmatrix} m_1 \cos^2 \theta + m_2 \sin^2 \theta & \frac{m_2 - m_1}{2} \sin(2\theta) \\ \frac{m_2 - m_1}{2} \sin(2\theta) & m_1 \sin^2 \theta + m_2 \cos^2 \theta \end{pmatrix}$$

Due to radiative correction this mass matrix gets modified in the low scale M_Z to [11–13]

$$\begin{aligned} M_F(M_Z) &= \begin{pmatrix} 1 + \delta_\alpha & 0 \\ 0 & 1 + \delta_\beta \end{pmatrix} M_F(\Lambda) \begin{pmatrix} 1 + \delta_\alpha & 0 \\ 0 & 1 + \delta_\beta \end{pmatrix} \\ &= \begin{pmatrix} (1 + 2\delta_\alpha)(m_1 \cos^2 \theta + m_2 \sin^2 \theta) & (1 + \delta_\alpha + \delta_\beta) \frac{m_2 - m_1}{2} \sin(2\theta) \\ (1 + \delta_\alpha + \delta_\beta) \frac{m_2 - m_1}{2} \sin(2\theta) & (1 + 2\delta_\beta)(m_1 \sin^2 \theta + m_2 \cos^2 \theta) \end{pmatrix} \\ &\equiv \begin{pmatrix} A & B \\ B & C \end{pmatrix} \end{aligned} \quad (1)$$

Here δ_α and δ_β are the radiative corrections to the flavours α and β .

The mixing angle at the low scale θ' in the limit of near degeneracy $m_1 \approx m_2 \approx m$ is given as

$$\begin{aligned} \tan 2\theta' &= \frac{2B}{C - A} \\ &= \frac{(1 + \delta_\alpha + \delta_\beta)(m_2 - m_1) \sin(2\theta)}{(m_2 - m_1) \cos(2\theta) + 2(\delta_\beta - \delta_\alpha)m} \end{aligned} \quad (2)$$

If

$$2(\delta_\beta - \delta_\alpha) = \frac{m_1 - m_2}{m} \cos(2\theta) \quad (3)$$

then $\theta' \rightarrow 45^\circ$

Let $\epsilon = 2(\delta_\beta - \delta_\alpha)$. Without loss of generality let $m_1 > m_2$. Whatever mixing angle θ one takes at high scale Λ , when one goes to low scale the mixing angle θ' will be maximal if condition (3) is satisfied [10]. We call this value of ϵ given by eq.(3) as ϵ_r (r referring to resonance). However it is crucial to investigate whether this condition can be satisfied at an

arbitrary mixing angle θ at the high scale. This is because the masses m_1 and m_2 in eq.(3) are the masses at the high scale Λ . Though the masses themselves don't have significant evolution through radiative correction the same cannot be said about the mass differences near degeneracy. To see this, consider the mass square difference at the low scale

$$\Delta m^2(M_Z) = (C + A)\sqrt{(C - A)^2 + 4B^2}$$

From eq.(2) this becomes

$$\Delta m^2(M_Z) \sin(2\theta') = 2B(C + A)$$

From eq.(1), in the limit of near degeneracy we get

$$\Delta m^2(M_Z) \sin(2\theta') = \Delta m^2(\Lambda) \sin(2\theta) \quad (4)$$

Here we have used the near degeneracy approximation in the following

$$\frac{m_2 - m_1}{m} = \frac{(m_2 - m_1)(m_2 + m_1)}{m(m_2 + m_1)} \approx \frac{\Delta m^2}{2m^2}$$

This is as expected in a MSW like scenario, where due to significant change in mass square difference, the mixing angle undergoes drastic enhancement in the resonance region. Eq.(4) says that once $\Delta m^2(M_Z)$ and $\sin(2\theta')$ are determined from oscillation experiments, the mass square difference and the mixing angle at the high scale Λ are related. So if the mixing angle θ is small at the high scale then the mass square difference $\Delta m^2(\Lambda)$ should be large. This would need a large ϵ to satisfy the resonance condition (3). We will see this in detail in the numerical plots below.

When θ' is maximal, we have from (4)

$$\Delta m^2(M_Z) = \Delta m^2(\Lambda) \sin(2\theta) \quad (5)$$

We see here that once the evolution of Δm^2 is fixed from the scale Λ to the scale M_Z , the mixing angle at the high scale Λ is fixed and not arbitrary.

Eq.(3) is a relation between ϵ_r and the mass difference and mixing angles at the high scale. It would be appropriate to get ϵ and ϵ_r in terms of the quantities at the low scale. Rearranging eq.(2) and using eq.(4) we get

$$\epsilon = \frac{\Delta m^2(M_Z)}{2m^2} \sin(2\theta') \left(\frac{1}{\tan(2\theta')} - \frac{1}{\tan(2\theta)} \right) \quad (6)$$

Since the combination $\Delta m^2(M_Z) \sin(2\theta')$ does not change under renormalisation group evolution as evident from eq.(4), we see from eq.(6) that given an ϵ if a small θ is enhanced to a large θ' around 45° then a large θ will end up at a θ' around 90° which is small mixing

but with inverted hierarchy. This observation is consistent with the explicit evolution of the neutrino mixing in SM, MSSM and two Higgs doublet model done in ref. [12] and [14].

ϵ_r is given by $\theta' = 45^\circ$ in eq.(6)

$$\epsilon_r = \frac{\Delta m^2(M_Z)}{2m^2} \left(-\frac{1}{\tan(2\theta)} \right) \quad (7)$$

From eq.(6) and (7) we get the range of ϵ that generates large mixing around the maximal. This is given as

$$\delta\epsilon = \epsilon_r - \epsilon \approx \frac{\Delta m^2(M_Z)}{2m^2} \left(\frac{\sin(2\theta') - 1}{\tan(2\theta)} \right) \quad (8)$$

where $\tan(2\theta') \gg \tan(2\theta)$ around $\theta' = \pi/4$ and small θ . So if we want the mixing at the low scale to be anything in the range of 30° to 45° the range of ϵ can be obtained by putting $\theta' = 30^\circ$ in eq.(8). This will be about 15% of ϵ_r . If we also consider the reversed hierarchy after maximal mixing, this range becomes 30%. With the current possibility of atmospheric neutrino mixing angle to be very near maximal ($40 - 50^\circ$), this range gets restricted to about 4% around ϵ_r .

While it may be possible for this mechanism to enhance some mixing angle from high scale to low scale with a few % range of ϵ around ϵ_r , another question we ask is, given an ϵ what is the range of mixing angle at the high scale that can be enhanced at the low scale. Figure (1) shows a plot of ϵ verses the high scale mixing angle θ at various values of the low scale mixing θ' as obtained from eq.(6). We have taken $\Delta m^2(M_Z) = 10^{-3}eV^2$ and $m = 1eV$. So this is the case for atmospheric neutrino solution. It is clear from the figure that for very small θ ($< 0.5^\circ$) the ϵ needed for the enhancement to occur is as large as 0.1. As ϵ is expected to be small, this mechanism cannot produce a large mixing from such small mixing at high scales. If $\epsilon \sim 0.01$ only θ in a small range $1.5^\circ - 2^\circ$ can be enhanced to near maximal mixing (30° to 45°) at the low scale. But if θ is over 2.5° then this mechanism becomes an overkill as the low scale mixing then again becomes small but with an inverted hierarchy. This is as expected in MSW mechanism when the neutrinos pass through the resonance and then go far away from it, the mixing is no longer maximal and the hierarchy gets inverted. If $\epsilon \sim 0.001$, then only θ between 10° to 15° get enhanced. While for $\epsilon = 0.0001$ high scale mixing upto about 25° don't get any significant enhancement. If the same analysis is done for the solution of solar neutrinos then the corresponding ϵ should be lowered by two orders of magnitude as $\Delta m^2(M_Z)$ is $\sim 10^{-5}eV^2$ instead of $10^{-3}eV^2$ in eq.(7). So we see that there is no range of ϵ that can enhance a large range of mixing angles at the high scale. Thus in order to generate maximal mixing at the low scale through radiative correction of the masses at the high scale, one has to start with specific mixing angle at the high scale depending upon the particular model one considers. Similar conclusion have been obtained in specific models [9,12,14] through renormalisation group evolution of neutrino mass matrix.

To summarise, we see that radiative correction cannot give a general enhancement of any mixing angles at the high scales to maximal mixing at the low scale. If it works due to fine tuning of parameters within a few percent for low mixing angles, it would kill the large mixing at high scales to small mixing at low scale. This is somewhat to be expected as large mixing between neutrinos of same CP parity is unstable near mass degeneracy [6–9]. Radiative magnification doesn't give the freedom to start with arbitrary mixing angle at high scale in building models of neutrino mass and mixing structures.

FIGURES




FIG. 1. ϵ required to generate θ' at the low scale as a function of θ at the high scale. The solid line, dotted line and the dashed lines are for $\theta' = 45^\circ, 30^\circ$ and 20° respectively. Here $\Delta m^2(M_Z) = 10^{-3} eV^2$.

REFERENCES

- [1] Y. Fukuda *et al*, Super-Kamiokande Collaboration, Phys. Rev. Lett. **81**, 1158 (1998) Erratum *ibid.* **81** 4279 (1998); *ibid* **82** 1810 (1999); Y. Suzuki, Nucl. Phys. Proc. Suppl. **77** 35 (1999); SAGE collaboration, J.N. Abdurashitov *et al*, Phys. Rev. **C60** 055801 (1991); GALLEX collaboration, W. Hampel *et al*, Phys. Lett. **B447** 127 (1999); SNO collaboration, Q. R. Ahmad *et al*, Phys. Rev. Lett. **87** 071301 (2001).
- [2] J.N. Bahcall, M.C. Gonzales-Garcia and C. Pena-Garay, JHEP **0108** 014 (2001); A. Bandopadhyay, S. Choubey, S. Goswami and K. Kar, Phys. Lett. **B519** 83 (2001); P. Krastev and A. Yu. Smirnov, hep-ph/0108177; G.L. Fogli, E. Lisi, D. Montanino and A. Palazzo, Phys. Rev. **D64** 093007 (2001); M.V. Garzelli and C. Giunti, JHEP **0112** 017 (2001); V. Barger, D. Marfatia and K. Whisnant, Phys. Rev. Lett. **88** 011302 (2002).
- [3] L. Wolfenstein, Phys. Lett. **107B** 77 (1981); Nucl. Phys. **B175**, 93 (1980); G.C. Branco, W. Grimus and L. Lavoura, Nucl. Phys. **B312** 492 (1989); S.T. Petcov and A. Yu. Smirnov, Phys. Lett. **B322** 109 (1994); Y.B. Zeldovich, Doklady Akad. Nauk. SSSR, **86** 505 (1952); E.J. Konopinski and H.M. Mahmoud, Phys. Rev. **92** 1045 (1953).
- [4] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, Proceedings of the workshop, Stony Brook, New York, 1979, edited by P. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1979),p. 315; T. Yanagida, Prog. Theor. Phys. **B135** 661 (1978).
- [5] G.Dutta and A.S. Joshipura, Phys. Rev. **D51** 3838 (1995).
- [6] E.Ma, J.Phys. **G25**, L97-L100 (1999)
- [7] N. Haba, Y. Matsui, N. Okamura, Eur.Phys.J. **C17** 513 (2000); N.Haba, Y.Matsui, N.Okamura, T.Suzuki, Phys.Lett. **B489** 184 (2000); N. Haba, Y. Matsui, N. Okamura, M. Sugiura, Prog.Theor.Phys. **103** 145 (2000); Eur.Phys.J. **C10** 677 (1999); N. Haba, N. Okamura, Eur.Phys.J. **C14** 347 (2000).
- [8] P.H. Chankowski and Stefan Pokorski, Int. J. Mod. Phys. **A17** 575 (2002); S. Antusch, M. Drees, J. Kersten, M. Lindner and M.Ratz, Phys. Lett. **B519**, 238 (2001).
- [9] J.A. Casas, J.R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. **B573**, 652 (2000).
- [10] K.R.S. Balaji, A.S. Dighe, R. N. Mohapatra, M.K.Parida, Phys. Rev. Lett.**84**, 5034 (2000)
- [11] J. Ellis and S. Lola, Phys. Lett. **B458**, 310 (1999).
- [12] K. S. Babu, C.N. Leung and J. Pantaleone, Phys. Lett. **B319**, 191 (1993);

- [13] P.H. Chankowski and Z. Pluciennik, Phys. Lett. **B316**, 312 (1993).
- [14] N. Haba, N. Okamura and M. Sugiura, Prog. Theor. Phys. **103**, 367 (2000).